

A39

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
1 May 2003 (01.05.2003)

PCT

(10) International Publication Number  
WO 03/035049 A2(51) International Patent Classification<sup>7</sup>: A61K 31/00

(21) International Application Number: PCT/IB02/04251

(22) International Filing Date:  
20 September 2002 (20.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/323,313 20 September 2001 (20.09.2001) US

(71) Applicant (for all designated States except US): AB SCIENCE [FR/FR]; 3, avenue George V, F-75008 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MOUSSY, Alain [FR/FR]; 22 bis, Passage Dauphine, F-75006 Paris (FR). KINET, Jean-Pierre [FR/US]; 3 Hunt Road, Lexington, MA 02421 (US).

(74) Agents: MARTIN, Jean-Jacques et al.; Cabinet Regimbeau, 20, rue de Chazelles, F-75847 Paris Cedex 17 (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

## Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF POTENT, SELECTIVE AND NON-TOXIC C-KIT INHIBITORS FOR TREATING BACTERIAL INFECTIONS

(57) Abstract: The present invention relates to a method for treating bacterial infections, preferably infections caused by FimH expressing bacteria, comprising administering a tyrosine kinase inhibitor to a human in need of such treatment, more particularly a non toxic, potent and selective c-kit inhibitor, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

WO 03/035049 A2

WO 03/035049

PCT/IB02/04251

I

**Use of potent, selective and non toxic c-kit inhibitors**  
**for treating bacterial infections**

5    The present invention relates to a method for treating bacterial infections, preferably infections caused by FimH expressing bacteria, comprising administering a tyrosine kinase inhibitor to a human in need of such treatment, more particularly a non toxic, potent and selective c-kit inhibitor, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

10

Bacterial infections are the most common diseases among mammalian and yet they remain deadly in case of resistant strains appearance. Resistance primarily originates from the extensive use of antibiotics. Antibiotics are agents acting on the bacterial cell wall such as bacitracin, the cephalosporins, and the penicillins, agents capable of  
15    inhibiting replication and protein synthesis by their effects on ribosomes, such as the aminoglycosides, the tetracyclines, the streptomycins and the macrolide antibiotics such as erythromycin; agents affecting nucleic acid metabolism, such as the fluoroquinolones, actinomycin; and drugs affecting intermediary metabolism, such as the sulfonamides and trimethoprim.

20

Despite the efficacy of antibiotics, few bacteria occasionally acquire mutations under high selection pressure, which renders the above mentioned antibiotic molecular targets insensitive and leads to the birth of new resistant strains.

More recently, multi-resistant strains have been observed during nosocomial infections  
25    and has come to the attention of the public. Facing the emergence of these deadly strains, research has focused on other mechanisms leading to multi-resistance. For example, it has been found that the marA loci confers multiple antibiotic resistance via increased

WO 03/035049

PCT/IB02/04251

2

efflux of many structurally unrelated antibiotics (McMurry et al., *Antimicrob. Agents Chemother.* 38:542-546, 1994). Multi-drug efflux pumps are now generally thought to be responsible for drugs insensitivity.

5 However, this mechanism leading to the resistance of bacteria does not explain the recurrence observed in bacterial infections. Indeed, after eradication of the bacteria, resurgence is observed later on suggesting that a small portion of bacteria were able to survive and remain concealed in the body. For example, urinary tract infections (UTI) have been treated for years with the antibiotics Bactrim, Macrochantin and a combination  
10 of Sulfa drugs that offer quick relief, but these antibiotics become useless after several prescriptions because the infection looks as if it has settled in the body and re-emerges from time to time.

Therefore, there is a need for new medications that would prevent and treat resurgence of  
15 bacterial infections.

In connection with the invention, it is postulated that bacteria, especially FimH expressing bacteria, are capable of escaping the immune system as well as the action of antibiotics by integration into mast cells, in which they remain concealed for a period  
20 time.

Mast cells (MC) are tissue elements derived from a particular subset of hematopoietic stem cells that express CD34, c-kit and CD13 antigens (Kirshenbaum et al, *Blood.* 94: 2333-2342, 1999 and Ishizaka et al, *Curr Opin Immunol.* 5: 937-43, 1993). Immature  
25 MC progenitors circulate in the bloodstream and differentiate in tissues. These differentiation and proliferation processes are under the influence of cytokines, one of

WO 03/035049

PCT/IB02/04251

3

utmost importance being Stem Cell Factor (SCF), also termed Kit ligand (KL), Steel factor (SL) or Mast Cell Growth Factor (MCGF). SCF receptor is encoded by the protooncogene c-kit, that belongs to type III receptor tyrosine kinase subfamily (Boissan and Arock, J Leukoc Biol. 67: 135-48, 2000). This receptor is also expressed on others

5 hematopoietic or non hematopoietic cells. Ligation of c-kit receptor by SCF induces its dimerization followed by its transphosphorylation, leading to the recruitment and activation of various intracytoplasmic substrates. These activated substrates induce multiple intracellular signaling pathways responsible for cell proliferation and activation (Boissan and Arock, 2000). Mast cells are characterized by their heterogeneity, not only

10 regarding tissue location and structure but also at the functional and histochemical levels (Aldenberg and Enerback., Histochem. J. 26: 587-96, 1994 ; Bradding et al. J Immunol. 155: 297-307, 1995 ; Irani et al, J Immunol. 147: 247-53, 1991 ; Miller et al, Curr Opin Immunol. 1: 637-42, 1989 and Welle et al, J Leukoc Biol. 61: 233-45, 1997).

15 Apart from their key role as effector cells of allergic and potentially lethal anaphylactic reactions, mast cells might contribute to the initiation of acquired immune reactions. Indeed, mast cells can phagocytosize diverse particles, and particularly bacteria. For example, recent studies have implicated rodent mast cells in the innate immune response to infectious bacteria and have shown that human mast cells are intrinsically

20 capable of mediating microbial recognition and of actively contributing to the host defense against bacteria ; Arock M et al, Infect Immun 1998 Dec;66(12):6030-4.

Galli SJ et al, Curr Opin Immunol 1999 Feb;11(1):53-9 suggested that mast cell function can be manipulated for therapeutic ends using SCF to boost immune response. This was also proposed by Maurer et al, J Exp Med 1998 Dec 21;188(12):2343-8 who identified

25 c-kit and mast cells as potential therapeutic targets for enhancing innate immune responses.

WO 03/035049

PCT/IB02/04251

4

While this can be acknowledged as far as acute infections are concerned, it could have serious drawbacks when considering recurrent bacterial infections.

Indeed, mast cells display very peculiar cell membrane structures called caveolae. Caveolae are subcellular structures implicated in the import and transcytosis of macromolecules and in transmembrane signaling. The composition and function of caveolae is reviewed in Anderson RG, Annu Rev Biochem 1998;67:199-225. In this article, caveolae are presented not just as an endocytic device with a peculiar membrane shape but rather as an entire membrane system with multiple functions essential for the cell. It is also mentioned that pathogens have been identified that use it as a means of gaining entrance to the cell.

Shin JS et al, Science. 2000 Aug 4;289(5480):732-3 reported that caveolae were detected in the microvilli and intracellular vesicles of cultured mouse bone marrow-derived mast cells (BMMCs). CD48, a receptor for FimH-expressing (type I fimbriated) *Escherichia coli*, was specifically localized to plasmalemmal caveolae in BMMCs. The involvement of caveolae in bacterial entry into BMMCs was demonstrated because caveolae-disrupting and -usurping agents specifically blocked *E. coli* entry. More importantly, it was demonstrated that some microbes utilize the unique features of caveolae to enter and traffic, without any apparent loss of viability and function, to different sites within immune and other host cells; Shin & Abraham, Immunology 2001, 102 (1), 2-7.

Therefore, bacteria-encapsulating caveolar chambers in mast cells form a reservoir of surviving bacteria that is postulated here to be implicated in the resurgence of infections.

WO 03/035049

PCT/IB02/04251

5

FimH, a mannose-binding lectin, expressed by many enterobacteria including *E. coli*, *K. pneumoniae* and *S. typhimurium*, binds to the receptor CD48 present at the surface of caveolae, Shin JS et al, FEMS Microbiol Lett 2001 Apr 13;197(2):131-8. As a result, FimH expressing bacteria enter inside mast cells and remain concealed and viable in  
5 caveolar chambers. In addition, Abraham SN et al, Nature 1988 Dec 15;336(6200):682-4 have observed a conservation of the D-mannose-adhesion protein among type 1 fimbriated members of the family Enterobacteriaceae.

It is proposed here that at some point, exocytosis of these chambers leads to the release  
10 of intact and living bacteria, which are responsible for the resurgence of the infection.

Consequently, apart from being beneficial for the organism through its ability to initiate immune responses towards a variety of pathogens, the mast cell may also be detrimental for the host during recurrent infectious diseases as specified above.

In such detrimental circumstances, therapeutic strategies aiming at blocking the  
15 activation and the survival of mast cells, for instance through inhibition of c-kit or c-kit signaling is proposed to decrease the inappropriate release of inflammatory mediators, as well as the survival of intracellular pathogens.

Therefore, the invention provides a new therapeutic strategy aimed at the use of c-kit specific kinase inhibitors to inhibit mast cell proliferation, survival and activation. A  
20 new route for treating recurrent bacterial infections is provided, which consists of destroying mast cells that constitute a reservoir for bacteria. It has been found that tyrosine kinase inhibitors and more particularly c-kit inhibitors are especially suited to reach this goal.

25

WO 03/035049

PCT/IB02/04251

6

**Description**

The present invention relates to a method for treating bacterial infections comprising administering a tyrosine kinase inhibitor to a mammalian in need of such treatment, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

Tyrosine kinase inhibitors are selected for example from bis monocyclic, bicyclic or heterocyclic aryl compounds (WO 92/20642), vinylene-azaindole derivatives (WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (US 5,330,992), Styryl compounds (US 5,217,999), styryl-substituted pyridyl compounds (US 5,302,606), seleoindoles and selenides (WO 94/03427), tricyclic polyhydroxylic compounds (WO 92/21660) and benzylphosphonic acid compounds (WO 91/15495), pyrimidine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones (US 5,792,783, EP 934 931, US 5,834,504, US 5,883,116, US 5,883,113, US 5,886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940) and aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758).

Preferably, said tyrosine kinase inhibitors are non-toxic, selective and potent c-kit inhibitors. Such inhibitors can be selected from the group consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives, quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic, bicyclic or heterocyclic aryl compounds, vinylene-azaindole derivatives and pyridyl-quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, , seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid compounds.


WO 03/035049

PCT/IB02/04251

7

Among preferred compounds, it is of interest to focus on pyrimidine derivatives such as N-phenyl-2-pyrimidine-amine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones (US 5,792,783, EP 934 931, US 5,834,504), US 5,883,116, US 5,883,113, US 5, 886,020, WO 96/40116 and WO  
5 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940), 4-amino-substituted quinazolines (US 3,470,182), 4-thienyl-2-(1H)-quinazolones, 6,7-dialkoxyquinazolines (US 3,800,039), aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO  
10 95/15758), 4-anilinoquinazoline compounds (US 4,464,375), and 4-thienyl-2-(1H)-quinazolones (US 3,551,427).

So, preferably, the invention relates to a method for treating bacterial infections comprising administering a non toxic, potent and selective c-kit inhibitor which is a  
15 pyrimidine derivative, more particularly N-phenyl-2-pyrimidine-amine derivatives of formula I :



wherein the R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>13</sub> to R<sub>17</sub> groups have the meanings depicted in EP 564 409  
20 B1, incorporated herein in the description.




WO 03/035049

PCT/IB02/04251

8

Preferably, the N-phenyl-2-pyrimidine-amine derivative is selected from the compounds corresponding to formula II :




5

Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;

R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;

- 10 and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function.

Preferably, R7 is the following group :



- 15 Among these compounds, the preferred are defined as follows:

R1 is a heterocyclic group, especially a pyridyl group,

R2 and R3 are H,

R4 is a C1-C3 alkyl, especially a methyl group,


R5 and R6 are H,

- 20 and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function, for example the group :

WO 03/035049


PCT/IB02/04251

9



Therefore, in a preferred embodiment, the invention relates to a method for treating bacterial infections comprising the administration of an effective amount of the compound known in the art as CGP57148B :

4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2-ylamino]phényl]-benzamide corresponding to the following formula :



The preparation of this compound is described in example 21 of EP 564 409 and the  $\beta$ -form, which is particularly useful is described in WO 99/03854.

Alternatively, the c-kit inhibitor can be selected from :

- indolinone derivatives, more particularly pyrrol-substituted indolinones,
- monocyclic, bicyclic aryl and heteroaryl compounds, quinazoline derivatives,
- and quinaxolines, such as 2-phényl-quinaxoline derivatives, for example 2-phenyl-6,7-dimethoxy quinaxoline.

In a preferred aspect, the invention contemplates the method mentioned above, wherein said c-kit inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

WO 03/035049

PCT/IB02/04251

10

The expression "bacterial infections" will be understood herein as recurrent bacterial infections, more particularly resurging infections after asymptomatic periods. Preferably, bacteria are FimH expressing bacteria such as Gram-negative enterobacteria which include but are not limited to well known pathogenic species such as *E. coli*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Citrobactor freundii* and *Salmonella typhimurium*. In connection with the invention, bacterial infections encompass recurrent urinary tract infections such as bacterial cystitis and respiratory tract infections.

10

In another embodiment, c-kit inhibitors as mentioned above are inhibitors of activated c-kit. In frame with the invention, the expression "activated c-kit" means a constitutively activated-mutant c-kit including at least one mutation selected from point mutations, deletions, insertions, but also modifications and alterations of the natural c-kit sequence (SEQ ID N°1). Such mutations, deletions, insertions, modifications and alterations can occur in the transphosphorylase domain, in the juxtamembrane domain as well as in any domain directly or indirectly responsible for c-kit activity. The expression "activated c-kit" also means herein SCF-activated c-kit. Preferred and optimal SCF concentrations for activating c-kit are comprised between  $5 \cdot 10^{-7}$  M and  $5 \cdot 10^{-6}$  M, preferably around  $2 \cdot 10^{-6}$  M. In a preferred embodiment, the activated-mutant c-kit in step a) has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants. In another preferred embodiment, the activated-mutant c-kit in step a) has a deletion in the juxtamembrane domain of c-kit. Such a deletion is for example between codon 573 and 579 called c-kit d(573-579). The point mutation V559G proximal to the juxtamembrane domain c-kit is also of interest.

25

WO 03/035049

PCT/IB02/04251

11

In this regard, the invention contemplates a method for treating bacterial infections comprising administering to a mammalian in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :

- 5 a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested; under conditions allowing the components (i) and (ii) to form a complex,  
b) selecting compounds that inhibit activated c-kit,  
c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

10

This screening method can further comprise the step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit (for example in the transphosphorylase domain), which are also capable of inhibiting SCF-activated c-kit wild.

- 15 Alternatively, in step a) activated c-kit is SCF-activated c-kit wild.

A best mode for practicing this method consists of testing putative inhibitors at a concentration above 10  $\mu$ M in step a). Relevant concentrations are for example 10, 15, 20, 25, 30, 35 or 40  $\mu$ M.

20

In step c), IL-3 is preferably present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

Examples of IL-3 dependent cells include but are not limited to :

- 25 - cell lines naturally expressing and depending on c-kit for growth and survival. Among such cells, human mast cell lines can be established using the following procedures :  
normal human mast cells can be infected by retroviral vectors containing sequences coding for a mutant c-kit comprising the c-kit signal peptide and a TAG sequence

WO 03/035049

PCT/IB02/04251

12

allowing to differentiate mutant c-kits from c-kit wild expressed in hematopoietic cells by means of antibodies.

This technique is advantageous because it does not induce cellular mortality and the genetic transfer is stable and gives satisfactory yields (around 20 %). Pure normal human  
5 mast cells can be routinely obtained by culturing precursor cells originating from blood obtained from human umbilical vein. In this regard, heparinated blood from umbilical vein is centrifuged on a Ficoll gradient so as to isolate mononucleated cells from other blood components. CD34+ precursor cells are then purified from the isolated cells mentioned above using the immunomagnetic selection system MACS (Miltenyi biotech).  
10 CD34+ cells are then cultured at 37°C in 5 % CO<sub>2</sub> atmosphere at a concentration of 10<sup>5</sup> cells per ml in the medium MCCM ( $\alpha$ -MEM supplemented with L-glutamine, penicillin, streptomycin, 5 10<sup>-5</sup> M  $\beta$ -mercaptoethanol, 20 % veal foetal serum, 1 % bovine albumin serum and 100 ng/ml recombinant human SCF. The medium is changed every 5 to 7 days. The percentage of mast cells present in the culture is assessed each week, using  
15 May-Grünwal Giemsa or Toluidine blue coloration. Anti-tryptase antibodies can also be used to detect mast cells in culture. After 10 weeks of culture, a pure cellular population of mast cells (< 98 %) is obtained.

It is possible using standard procedures to prepare vectors expressing c-kit for transfecting the cell lines established as mentioned above. The cDNA of human c-kit has  
20 been described in Yarden et al., (1987) EMBO J.6 (11), 3341-3351. The coding part of c-kit (3000 bp) can be amplified by PCR and cloned, using the following oligonucleotides :

- 5'AAGAAGAGATGGTACCTCGAGGGGTGACCC3' (SEQ ID No2) sens
  - 5'CTGCTTCGCGGCCGCGTAACTCTTCTCAACCA3' (SEQ ID No3)
- 25 antisens

WO 03/035049

PCT/IB02/04251

13

The PCR products, digested with NotI and XhoI, has been inserted using T4 ligase in the pFlag-CMV vector (SIGMA), which vector is digested with NotI and XhoI and dephosphorylated using CIP (Biolabs). The pFlag-CMV-c-kit is used to transform bacterial clone XLI-blue. The transformation of clones is verified using the following

5 primers :

- 5'AGCTCGTTTAGTGAACCGTC3' (SEQ ID No4) sens,
- 5'GTCAGACAAAATGATGCAAC3' (SEQ ID No5) antisens.

Directed mutagenesis is performed using relevant cassettes is performed with routine and common procedure known in the art..

- 10 The vector Migr-1 (ABC) can be used as a basis for constructing retroviral vectors used for transfecting mature mast cells. This vector is advantageous because it contains the sequence coding for GFP at the 3' and of an IRES. These features allow to select cells infected by the retrovirus using direct analysis with a fluorocytometer. As mentioned above, the N-terminal sequence of c-kit c-DNA can be modified so as to introduce a Flag
- 15 sequence that will be useful to discriminating heterogeneous from endogenous c-kit.

Other IL-3 dependent cell lines that can be used include but are not limited to:

- BaF3 mouse cells expressing wild-type or mutated form of c-kit (in the juxtamembrane and in the catalytic sites) are described in Kitayama et al, (1996), Blood
- 20 88, 995-1004 and Tsujimura et al, (1999), Blood 93, 1319-1329.
- IC-2 mouse cells expressing either c-kit<sup>WT</sup> or c-kit<sup>D814Y</sup> are presented in Piao et al, (1996), Proc. Natl. Acad. Sci. USA 93, 14665-14669.

IL-3 independent cell lines are :

WO 03/035049

PCT/IB02/04251

14

- HMC-1, a factor-independent cell line derived from a patient with mast cell leukemia, expresses a juxtamembrane mutant c-kit polypeptide that has constitutive kinase activity (Furitsu T et al, J Clin Invest. 1993;92:1736-1744 ; Butterfield et al, Establishment of an immature mast cell line from a patient with mast cell leukemia. Leuk Res. 1988;12:345-  
5 355 and Nagata et al, Proc Natl Acad Sci U S A. 1995;92:10560-10564).
- P815 cell line (mastocytoma naturally expressing c-kit mutation at the 814 position) has been described in Tsujimura et al, (1994), Blood 83, 2619-2626.

10 The extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*. In case it is measured *in vivo*, cell lines expressing an activated-mutant c-kit, which has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, are preferred.

15 Example of cell lines expressing an activated-mutant c-kit are as mentioned above.

In another preferred embodiment, the method further comprises the step consisting of testing and selecting compounds capable of inhibiting c-kit wild at concentration below 1  $\mu$ M. This can be measured *in vitro* or *in vivo*.

20 Therefore, compounds are identified and selected according to the method described above are potent, selective and non-toxic c-kit wild inhibitors.

Alternatively, the screening method according to the invention can be practiced *in vitro*. In this regard, the inhibition of mutant-activated c-kit and/or c-kit wild can be measured  
25 using standard biochemical techniques such as immunoprecipitation and western blot. Preferably, the amount of c-kit phosphorylation is measured.

WO 03/035049

PCT/IB02/04251

15

In a still further embodiment, the invention contemplates a method for treating bacterial infections as depicted above wherein the screening comprises :

- a) performing a proliferation assay with cells expressing a mutant c-kit (for example in the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an  $IC_{50} < 10 \mu M$ , by measuring the extent of cell death,
- b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,
- c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an  $IC_{50} < 10 \mu M$ , preferably an  $IC_{50} < 1 \mu M$ , by measuring the extent of cell death.

15

Here, the extent of cell death can be measured by  $^3H$  thymidine incorporation, the trypan blue exclusion method or flow cytometry with propidium iodide. These are common techniques routinely practiced in the art.

- Therefore, the invention embraces the use of the compounds defined above to manufacture a medicament for treating bacterial infections in mammalian, especially in human. Such medicament is particularly useful for the treatment of recurrent bacterial infections, more particularly resurging infections after asymptomatic periods such as bacterial cystitis and respiratory tract infections. Preferably, the invention contemplates the use of the compounds defined above to manufacture a medicament for treating FimH expressing bacteria infections such as Gram-negative enterobacteria which include but are not limited to *E. coli*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Citrobacter freundii* and *Salmonella typhimurium*.



WO 03/035049

PCT/IB02/04251

16

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, 5 intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used 10 pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, Pa.).

Pharmaceutical compositions for oral administration can be formulated using 15 pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

20 Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and 25 proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

WO 03/035049

PCT/IB02/04251

17

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or  
5 dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or  
10 sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

15

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such  
20 as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain  
25 suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

WO 03/035049

PCT/IB02/04251

18

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succine, acids, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred  
5 preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

Pharmaceutical compositions suitable for use in the invention include compositions  
10 wherein c-kit inhibitors are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or  
15 experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. As mentioned above, a tyrosine kinase inhibitor and more particularly a c-kit  
20 inhibitor according to the invention is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

In another embodiment, the invention is aimed at a product comprising a tyrosine kinase inhibitor, more particularly a c-kit inhibitor, and at least one antibiotic selected  
25 bacitracin, the cephalosporins, the penicillins, the aminoglycosides, the tetracyclines, the streptomycins and the macrolide antibiotics such as erythromycin; the fluoroquinolones, actinomycin, the sulfonamides and trimethoprim for a separate, sequential or

WO 03/035049

PCT/IB02/04251

19

simultaneous use for treating recurrent bacterial infections, resurging infections after asymptomatic periods such as bacterial cystitis and respiratory tract infections.

This product is particularly useful for treating FimH expressing bacteria infections such as Gram-negative enterobacteria including *E. coli*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Citrobacter freundii* and *Salmonella typhimurium*. Preferably, said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3 and the product further comprises an acceptable pharmaceutical carrier suitable for oral administration.

10

WO 03/035049

PCT/IB02/04251

20

## CLAIMS


1. A method for treating bacterial infections comprising administering a tyrosine kinase  
5 inhibitor to a mammalian in need of such treatment, wherein said inhibitor is unable to  
promote death of IL-3 dependent cells cultured in presence of IL-3.
2. A method according to claim 1, wherein said tyrosine kinase inhibitor is a non-toxic,  
selective and potent c-kit inhibitor.
- 10 3. A method according to claim 2, wherein said inhibitor is selected from the group  
consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives,  
quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic,  
bicyclic or heterocyclic aryl compounds, vinylene-azaindole derivatives and pyridyl-  
15 quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, ,  
seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid  
compounds.
4. A method for treating bacterial infections comprising administering a non toxic,  
20 potent and selective c-kit inhibitor to a mammalian in need of such treatment, selected  
from the group consisting of :
- pyrimidine derivatives, more particularly N-phenyl-2-pyrimidine-amine derivatives.
  - indolinone derivatives, more particularly pyrrol-substituted indolinones,
  - monocyclic, bicyclic aryl and heteroaryl compounds,
  - 25 - and quinazoline derivatives.

WO 03/035049

PCT/IB02/04251

21


5. A method according to claim 2, wherein said inhibitor is selected from the group consisting of N-phenyl-2-pyrimidine-amine derivatives having the formula II :



Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;

R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function, preferably the following group :



6. A method according to claim 5, wherein said inhibitor is the 4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2 ylamino]phényl]-benzamide.

7. A method according to one of claims 2 to 6, wherein said c-kit inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

20

WO 03/035049

PCT/IB02/04251

22

8. A method according to one of claims 2 to 7, wherein said inhibitor is an inhibitor of activated c-kit selected from a constitutively activated-mutant c-kit and/or SCF-activated c-kit.
- 5 9. A method according to claim 8, wherein the activated-mutant c-kit has at least one mutation selected from mutations proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, and a deletion in the juxtamembrane domain of c-kit, preferably between codon 573 and 579.
- 10 10. A method for treating bacterial infections comprising administering to a mammalian in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :
- 15 a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested;  
under conditions allowing the components (i) and (ii) to form a complex,  
b) selecting compounds that inhibit activated c-kit,  
c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.
- 20 11. A method according to claim 10, wherein the screening method further comprises the step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit, which are also capable of inhibiting SCF-activated c-kit wild.
- 25 12. A method according to claim 10, wherein activated c-kit is SCF-activated c-kit wild.
13. A method according to one of claims 10 to 12, wherein putative inhibitors are tested at a concentration above 10  $\mu$ M in step a).

WO 03/035049

PCT/IB02/04251

23

14. A method according to one of claims 10 to 13, wherein IL-3 is present in the culture media of IL-3 dependent cells at a concentration comprised between between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

5 15. A method according to one of claims 10 to 14, wherein the extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*.

16. A method according to one of claims 10 to 15, wherein the screening method further comprises the step consisting of testing and selecting *in vitro* or *in vivo* compounds  
10 capable of inhibiting c-kit wild at concentration below 1  $\mu$ M.

17. A method according to claim 16, wherein the test is performed using cells lines selected from the group consisting of mast cells, transfected mast cells, BaF3, and IC-2.

15 18. A method according to claim 16, wherein the test includes the determination of the amount of c-kit phosphorylation.

19. A method for treating bacterial infections according to one of claims 10 to 18, wherein the screening comprises :

- 20 a) performing a proliferation assay with cells expressing a mutant c-kit (for example in the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an  $IC_{50} < 10 \mu M$ , by measuring the extent of cell death,
- b) performing a proliferation assay with cells expressing c-kit wild said subset of  
25 candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,



WO 03/035049

PCT/IB02/04251

24

c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an  $IC_{50} < 10 \mu M$ , preferably an  $IC_{50} < 1 \mu M$ , by measuring the extent of cell death.

5

20. A method according to one of claims 1 to 19 for treating recurrent bacterial infections, more particularly resurging infections after an asymptomatic period.

21. A method according to one of claims 1 to 19 for treating bacterial infections, wherein  
10 bacteria are FimH expressing bacteria such as Gram-negative enterobacteria such as *E. coli*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Citrobactor freundii* and *Salmonella typhimurium*.

22. A method according to one of claims 1 to 19 for treating urinary tract infections such  
15 as bacterial cystitis and respiratory tract infections.

23. A method according to one of claims 20 to 22, wherein the inhibitor is administered orally.

20 24. Use of a tyrosine kinase inhibitor, more particularly a c-kit inhibitor, to manufacture a medicament for treating bacterial infections in mammalian, especially in human, preferably for the treatment of recurrent bacterial infections, resurging infections after asymptomatic periods such as bacterial cystitis and respiratory tract infections.

25 25. Use of a tyrosine kinase inhibitor, more particularly a c-kit inhibitor, to manufacture a medicament for treating FimH expressing bacteria infections such as Gram-negative enterobacteria including *E. coli*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Citrobactor freundii* and *Salmonella typhimurium*.

WO 03/035049

PCT/IB02/04251

25

26. A product comprising a tyrosine kinase inhibitor, more particularly a c-kit inhibitor, and at least one antibiotic selected from bacitracin, the cephalosporins, the penicillins, the aminoglycosides, the tetracyclines, the streptomycins and the macrolide antibiotics  
5 such as erythromycin; the fluoroquinolones, actinomycin, the sulfonamides and trimethoprim for a separate, sequential or simultaneous use for treating recurrent bacterial infections, resurging infections after asymptomatic periods such as bacterial cystitis and respiratory tract infections.

10 27. A product according to claim 26 for treating FimH expressing bacteria infections such as Gram-negative enterobacteria including *E. coli*, *Klebsiella pneumoniae*, *Serratia marcescens*, *Citrobacter freundii* and *Salmonella typhimurium*.

28. A product according to one of claims 26 and 27, wherein said c-kit inhibitor is  
15 unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

29. A product according to one of claims 26 and 27, further comprising an acceptable pharmaceutical carrier suitable for oral administration.

20

WO 03/035049

1/4

PCT/IB02/04251

## SEQUENCE LISTING

&lt;110&gt; AB Science

<120> Use of potent, selective and non toxic c-kit inhibitors  
for treating bacterial infections

&lt;130&gt; D19831 NT

&lt;150&gt; US 60/323,313

&lt;151&gt; 2001-09-20

&lt;160&gt; 5

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 976

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;223&gt; Human c-kit

&lt;400&gt; 1

Met Arg Gly Ala Arg Gly Ala Trp Asp Phe Leu Cys Val Leu Leu Leu  
1 5 10 15Leu Leu Arg Val Gln Thr Gly Ser Ser Gln Pro Ser Val Ser Pro Gly  
20 25 30Glu Pro Ser Pro Pro Ser Ile His Pro Gly Lys Ser Asp Leu Ile Val  
35 40 45Arg Val Gly Asp Glu Ile Arg Leu Leu Cys Thr Asp Pro Gly Phe Val  
50 55 60Lys Trp Thr Phe Glu Ile Leu Asp Glu Thr Asn Glu Asn Lys Gln Asn  
65 70 75 80Glu Trp Ile Thr Glu Lys Ala Glu Ala Thr Asn Thr Gly Lys Tyr Thr  
85 90 95Cys Thr Asn Lys His Gly Leu Ser Asn Ser Ile Tyr Val Phe Val Arg  
100 105 110Asp Pro Ala Lys Leu Phe Leu Val Asp Arg Ser Leu Tyr Gly Lys Glu  
115 120 125Asp Asn Asp Thr Leu Val Arg Cys Pro Leu Thr Asp Pro Glu Val Thr  
130 135 140Asn Tyr Ser Leu Lys Gly Cys Gln Gly Lys Pro Leu Pro Lys Asp Leu  
145 150 155 160Arg Phe Ile Pro Asp Pro Lys Ala Gly Ile Met Ile Lys Ser Val Lys  
165 170 175Arg Ala Tyr His Arg Leu Cys Leu His Cys Ser Val Asp Gln Glu Gly  
180 185 190Lys Ser Val Leu Ser Glu Lys Phe Ile Leu Lys Val Arg Pro Ala Phe  
195 200 205Lys Ala Val Pro Val Val Ser Val Ser Lys Ala Ser Tyr Leu Leu Arg  
210 215 220

WO 03/035049

2/4

PCT/IB02/04251

Glu	Gly	Glu	Glu	Phe	Thr	Val	Thr	Cys	Thr	Ile	Lys	Asp	Val	Ser	Ser	225	230	235	240
Ser	Val	Tyr	Ser	Thr	Trp	Lys	Arg	Glu	Asn	Ser	Gln	Thr	Lys	Leu	Gln	245	250	255	
Glu	Lys	Tyr	Asn	Ser	Trp	His	His	Gly	Asp	Phe	Asn	Tyr	Glu	Arg	Gln	260	265	270	
Ala	Thr	Leu	Thr	Ile	Ser	Ser	Ala	Arg	Val	Asn	Asp	Ser	Gly	Val	Phe	275	280	285	
Met	Cys	Tyr	Ala	Asn	Asn	Thr	Phe	Gly	Ser	Ala	Asn	Val	Thr	Thr	Thr	290	295	300	
Leu	Glu	Val	Val	Asp	Lys	Gly	Phe	Ile	Asn	Ile	Phe	Pro	Met	Ile	Asn	305	310	315	320
Thr	Thr	Val	Phe	Val	Asn	Asp	Gly	Glu	Asn	Val	Asp	Leu	Ile	Val	Glu	325	330	335	
Tyr	Glu	Ala	Phe	Pro	Lys	Pro	Glu	His	Gln	Gln	Trp	Ile	Tyr	Met	Asn	340	345	350	
Arg	Thr	Phe	Thr	Asp	Lys	Trp	Glu	Asp	Tyr	Pro	Lys	Ser	Glu	Asn	Glu	355	360	365	
Ser	Asn	Ile	Arg	Tyr	Val	Ser	Glu	Leu	His	Leu	Thr	Arg	Leu	Lys	Gly	370	375	380	
Thr	Glu	Gly	Gly	Thr	Tyr	Thr	Phe	Leu	Val	Ser	Asn	Ser	Asp	Val	Asn	385	390	395	400
Ala	Ala	Ile	Ala	Phe	Asn	Val	Tyr	Val	Asn	Thr	Lys	Pro	Glu	Ile	Leu	405	410	415	
Thr	Tyr	Asp	Arg	Leu	Val	Asn	Gly	Met	Leu	Gln	Cys	Val	Ala	Ala	Gly	420	425	430	
Phe	Pro	Glu	Pro	Thr	Ile	Asp	Trp	Tyr	Phe	Cys	Pro	Gly	Thr	Glu	Gln	435	440	445	
Arg	Cys	Ser	Ala	Ser	Val	Leu	Pro	Val	Asp	Val	Gln	Thr	Leu	Asn	Ser	450	455	460	
Ser	Gly	Pro	Pro	Phe	Gly	Lys	Leu	Val	Val	Gln	Ser	Ser	Ile	Asp	Ser	465	470	475	480
Ser	Ala	Phe	Lys	His	Asn	Gly	Thr	Val	Glu	Cys	Lys	Ala	Tyr	Asn	Asp	485	490	495	
Val	Gly	Lys	Thr	Ser	Ala	Tyr	Phe	Asn	Phe	Ala	Phe	Lys	Gly	Asn	Asn	500	505	510	
Lys	Glu	Gln	Ile	His	Pro	His	Thr	Leu	Phe	Thr	Pro	Leu	Leu	Ile	Gly	515	520	525	
Phe	Val	Ile	Val	Ala	Gly	Met	Met	Cys	Ile	Ile	Val	Met	Ile	Leu	Thr	530	535	540	
Tyr	Lys	Tyr	Leu	Gln	Lys	Pro	Met	Tyr	Glu	Val	Gln	Trp	Lys	Val	Val	545	550	555	560
Glu	Glu	Ile	Asn	Gly	Asn	Asn	Tyr	Val	Tyr	Ile	Asp	Pro	Thr	Gln	Leu	565	570	575	

WO 03/035049

PCT/IB02/04251

3/4

Pro Tyr Asp His Lys Trp Glu Phe Pro Arg Asn Arg Leu Ser Phe Gly  
 580 585 590  
 Lys Thr Leu Gly Ala Gly Ala Phe Gly Lys Val Val Glu Ala Thr Ala  
 595 600 605  
 Tyr Gly Leu Ile Lys Ser Asp Ala Ala Met Thr Val Ala Val Lys Met  
 610 615 620  
 Leu Lys Pro Ser Ala His Leu Thr Glu Arg Glu Ala Leu Met Ser Glu  
 625 630 635 640  
 Leu Lys Val Leu Ser Tyr Leu Gly Asn His Met Asn Ile Val Asn Leu  
 645 650 655  
 Leu Gly Ala Cys Thr Ile Gly Gly Pro Thr Leu Val Ile Thr Glu Tyr  
 660 665 670  
 Cys Cys Tyr Gly Asp Leu Leu Asn Phe Leu Arg Arg Lys Arg Asp Ser  
 675 680 685  
 Phe Ile Cys Ser Lys Gln Glu Asp His Ala Glu Ala Ala Leu Tyr Lys  
 690 695 700  
 Asn Leu Leu His Ser Lys Glu Ser Ser Cys Ser Asp Ser Thr Asn Glu  
 705 710 715 720  
 Tyr Met Asp Met Lys Pro Gly Val Ser Tyr Val Val Pro Thr Lys Ala  
 725 730 735  
 Asp Lys Arg Arg Ser Val Arg Ile Gly Ser Tyr Ile Glu Arg Asp Val  
 740 745 750  
 Thr Pro Ala Ile Met Glu Asp Asp Glu Leu Ala Leu Asp Leu Glu Asp  
 755 760 765  
 Leu Leu Ser Phe Ser Tyr Gln Val Ala Lys Gly Met Ala Phe Leu Ala  
 770 775 780  
 Ser Lys Asn Cys Ile His Arg Asp Leu Ala Ala Arg Asn Ile Leu Leu  
 785 790 795 800  
 Thr His Gly Arg Ile Thr Lys Ile Cys Asp Phe Gly Leu Ala Arg Asp  
 805 810 815  
 Ile Lys Asn Asp Ser Asn Tyr Val Val Lys Gly Asn Ala Arg Leu Pro  
 820 825 830  
 Val Lys Trp Met Ala Pro Glu Ser Ile Phe Asn Cys Val Tyr Thr Phe  
 835 840 845  
 Glu Ser Asp Val Trp Ser Tyr Gly Ile Phe Leu Trp Glu Leu Phe Ser  
 850 855 860  
 Leu Gly Ser Ser Pro Tyr Pro Gly Met Pro Val Asp Ser Lys Phe Tyr  
 865 870 875 880  
 Lys Met Ile Lys Glu Gly Phe Arg Met Leu Ser Pro Glu His Ala Pro  
 885 890 895  
 Ala Glu Met Tyr Asp Ile Met Lys Thr Cys Trp Asp Ala Asp Pro Leu  
 900 905 910  
 Lys Arg Pro Thr Phe Lys Gln Ile Val Gln Leu Ile Glu Lys Gln Ile  
 915 920 925

WO 03/035049

4/4

PCT/IB02/04251

Ser Glu Ser Thr Asn His Ile Tyr Ser Asn Leu Ala Asn Cys Ser Pro  
 930 935 940

Asn Arg Gln Lys Pro Val Val Asp His Ser Val Arg Ile Asn Ser Val  
 945 950 955 960

Gly Ser Thr Ala Ser Ser Ser Gln Pro Leu Leu Val His Asp Asp Val  
 965 970 975

<210> 2  
 <211> 30  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <223> Primer

<400> 2  
 aagaagagat ggtacctcga ggggtgaccc

30

<210> 3  
 <211> 33  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <223> Primer

<400> 3  
 ctgcttcgcg gccgcgttaa ctcttctcaa cca

33

<210> 4  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <223> Primer

<400> 4  
 agctcgttta gtgaaccgtc

20

<210> 5  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <223> Primer

<400> 5  
 gtcagacaaa atgatgcaac

20